

Further Reflections on Thompson's Conjecture*

Guizun Chen

ORE

ed by Elsevier - Publisher Connector

People's Republic of China 400715

Communicated by Walter Feit

Received October 22, 1996

Let G be a finite group and let $N(G) = \{n \in N \mid G \text{ has a conjugacy class } C, \text{ such that } |C| = n\}$. Professor J. G. Thompson has conjectured that: If G is a finite group with $Z(G) = 1$ and M a non-abelian simple group satisfying $N(G) = N(M)$, then $G \cong M$.

We have proved previously that: If M is a sporadic simple group or a simple group having its prime graph with at least three prime graph components, then Thompson's conjecture is correct. In this paper, we shall prove:

MAIN THEOREM. Let G be a finite group with $Z(G) = 1$ and $M = G_2(q)$ or $G_2(2)'$, where $q \geq 2$, such that $N(G) = N(M)$. Then $G \cong M$. © 1999 Academic Press

Key Words: Finite groups; conjugacy classes; characterization of a finite simple group

All notation are the same as in [4].

The next two lemmas follow from Lemmas 1.4 and 1.5 in [4].

LEMMA 1. Suppose G and M are two finite groups satisfying $t(M) \geq 2$, $N(G) = N(M)$, and $Z(G) = 1$. Then $|G| = |M|$.

LEMMA 2. Let G_1 and G_2 be finite groups satisfying $|G_1| = |G_2|$ and $N(G_1) = N(G_2)$. Then $t(G_1) = t(G_2)$ and $T(G_1) = T(G_2)$.

The next lemma follows from Theorem 2 in [2].

LEMMA 3. Suppose G is a Frobenius group of even order and H and K are the Frobenius kernel and the Frobenius complement of G , respectively. Then

* This work was partially supported by NNSF of China and Ph.D.F. of Southwest China Normal University.



$t(G) = 2$, $T(G) = \{\pi(H), \pi(K)\}$, and G has one of the following structures:

- (1) $2 \in \pi(H)$, all Sylow subgroups of K are cyclic.
- (2) $2 \in \pi(K)$, H is an abelian group, K is a solvable group, the Sylow subgroups of odd order of G are cyclic groups, and the 2-Sylow subgroups of G are cyclic or generalized quaternion groups.
- (3) $2 \in \pi(K)$, H is an abelian group, and there exists $K_0 \leq K$ such that $[K : K_0] \leq 2$, $K_0 = \mathbf{Z} \times SL(2, 5)$, $(|Z|, 2 \times 3 \times 5) = 1$, and the Sylow subgroups of \mathbf{Z} are cyclic.

The next lemma follows from Theorem 2 in [2].

LEMMA 4. Let G be a 2-Frobenius group of even order. Then $t(G) \geq 2$ and G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that $\pi(K/H) = \pi_2$, $\pi(H) \cup \pi(G/K) = \pi_1$, $|G/K| \mid |\text{Aut}(K/H)|$, and G/K and K/H are cyclic. Especially, $|G/K| < |K/H|$, and G is solvable.

LEMMA 5. If G is a finite group with more than one prime graph component, then G is one of the following: (a) Frobenius or 2-Frobenius, (b) simple, (c) an extension of a π_1 -group by a simple group, (d) simple by π_1 -solvable, or (e) a π_1 -group by simple by π_1 -group, where π_1 is the prime graph component containing prime 2 (see [8]).

LEMMA 6. If G is a finite group with more than one prime graph component and has a normal series $H \trianglelefteq K \trianglelefteq G$ such that H and G/K are π_1 -groups and K/H is simple, then H is a nilpotent group (see Lemma 3 in [8]).

LEMMA 7. Let G be a finite group with $Z(G) = 1$. If M is a non-abelian simple group with $t(M) = 2$ satisfying $N(G) = N(M)$, then

- (1) $|G| = |M|$, $t(G) = t(M)$, and $T(G) = T(M)$.
- (2) Let $|M| = m_1 m_2$, $T(M) = \{\pi_1, \pi_2\}$, and $\pi(m_i) = \pi_i$ for $i = 1$ or 2. Then $|G| = m_1 m_2$ and one of the following holds:
 - (a) G is a Frobenius or 2-Frobenius group;
 - (b) G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that G/K is a π_1 -group, H is a nilpotent π_1 -group, and K/H is a non-abelian simple group. Moreover $T(K/H) = \{\pi'_1, \pi'_2, \dots, \pi'_s, \pi_2\}$, $|K/H| = m'_1 m'_2 \cdots m'_s m_2$ and $m'_1 m'_2 \cdots m'_s \mid m_1$ where $\pi(m'_j) = \pi'_j$, $1 \leq j \leq s$.
 - (3) $|G/K| \mid |\text{Out}(K/H)|$.

Proof. (1) and (2) follow from Lemmas 1–6. Since $t(G) \geq 2$, we have $t(G/H) \geq 2$. Otherwise $t(G/H) = 1$, so that $t(G) = 1$ since $2 \mid |H|$ and H

a π_1 -group, a contradiction. Moreover we have $Z(G/H) = 1$. For any $xH \in G/H$ and $xH \notin K/H$, xH induces an automorphism of K/H . And this automorphism is trivial iff $xH \in Z(G/H)$. Therefore $G/K \leq \text{Out}(K/H)$ for $Z(G/H) = 1$, from which (3) follows.

Let G be a finite group of even order and $T(G) = \{\pi_1, \pi_2, \dots, \pi_t\}$, while $2 \in \pi_1$. Then $|G| = m_1 m_2 \cdots m_t$ such that $\pi(m_i) = \pi_i$, $i = 1, 2, \dots, t$. These m_i are called order components of G in [3, 4]. We use $\text{OC}(G)$ to denote the set of order components of G . And by [6, 8] we can easily calculate the order components of simple groups M with $t(M) \geq 2$ (see Tables 1–4 in [3]).

LEMMA 8. *Let G be a finite group with $t(\Gamma(G)) \geq 2$ and N a normal subgroup of G . If N is a π_i -group for some prime graph component π_i of G and a_1, a_2, \dots, a_r are some order components of G but not a π_i number, then $a_1 a_2 \cdots a_r$ is a divisor of $|N| - 1$.*

Proof. For any $x \in N$ and $x \neq 1$, the centralizer of x in G must be a π_i -group and so a_1, a_2, \dots, a_r divides $|\text{Cl}_G(x)|$. Therefore $a_1 a_2 \cdots a_r$ divides $|N| - 1$ for $N \trianglelefteq G$ and N is a union of conjugacy classes.

In the following lemma we use $D(G)$ to denote the odd order component of G . For $M = G_2(q)$ we have $D(M) = q^2 + q + 1$ for $3|q + 1$ and $q^2 - q + 1$ for $3|q - 1$.

LEMMA 9. *Let $M = G_2(q)$, $q \geq 4$, $q \equiv -1$ or $1 \pmod{3}$. Let $S_p \in \text{Syl}_p(M)$, where $p \in \pi(M)$ and $p \nmid q$. Then*

$$(1) \quad |S_p| < q^3 \text{ or } |S_p| = p^6.$$

(2) *If $p \in \pi_1(M)$ and $p^\alpha \mid |M|$, then $p^\alpha - 1 \equiv 0 \pmod{D(M)}$ if and only if $p|q$ or $q = 4$, $p^\alpha = 27$ or $q = 5$, $p^\alpha = 32$, where $\pi_1(M)$ is the prime graph component containing 2.*

Proof. Since $|G| = q^6(q^2 - q + 1)(q^2 + q + 1)(q - 1)^2(q + 1)^2$, it is easy to show that (1) holds.

To prove (2), it is sufficient to prove the necessity. And in the rest of the proof, we assume that $p \nmid q$. Since $p^\alpha - 1 \equiv 0 \pmod{D(M)}$, $p^\alpha > D(M)$.

Case 1. If $q \equiv 1 \pmod{3}$, then p^α is a factor of $4(q - 1)^2$, $4(q + 1)^2$, $3(q - 1)^2$, and $(q^2 + q + 1)/3$.

(a) If $p^\alpha | 4(q - 1)^2$, then, since $3|q - 1$, we have $p^\alpha \mid [(q - 1)^2 \text{ or } p^\alpha \mid 4(q - 1)^2/9]$. Thus $p^\alpha < q^2 - q + 1 = D(G)$, a contradiction.

(b) Suppose $p^\alpha | 4(q + 1)^2$.

First we further assume that $p = 2$. If $q + 1$ is not a power of 2, then, since $3 \nmid q + 1$, $p^\alpha \leq 4(q + 1)^2/25 < D(G)$, a contradiction. If $q + 1$ is a

power of 2, then $p^\alpha = (q+1)^2$, $2(q+1)^2$, or $4(q+1)^2$ for $p^\alpha > D(G)$ and $(q+1)^2/2 < D(M)$.

If $p^\alpha = (q+1)^2$, then, since $(q+1)^2 - 1 = D(G) + 2q - 1$ and $p^\alpha - 1 \equiv 0 \pmod{D(G)}$, we have $D(G) = 2q - 1$. Thus $q = 2$, a contradiction.

If $p^\alpha = 2(q+1)^2$, then $p^\alpha - 1 = 2D(G) + 6q - 1$. It is easy to check that this is impossible by $D(G) \mid p^\alpha - 1$. Similarly, it can be proved that $p^\alpha = 4(q+1)^2$ is also impossible.

We can deal with the case $p \neq 2$ similarly.

(c) If $p^\alpha \mid (q^2 + q + 1)/3$, then $P^\alpha < D(M)$, a contradiction.

(d) If $p^\alpha \mid 3(q-1)^2$, then $p^\alpha - 1 = 3D(M) - 3q - 1$, which means $3q + 1 = q^2 - q + 1$ and so $q = 4$ and $p^\alpha = 3^3$.

Case 2. If $q \equiv -1 \pmod{3}$, then p^α is a factor of $4(q-1)^2$, $4(q+1)^2$, $3(q+1)^2$, and $(q^2 - q + 1)/3$. Using the same method as in Case 1, we see that $q = 5$ and $p^\alpha = 32$.

By Tables 2 and 4 in [4] we know that if $3 \mid q$, then $\text{OC}(G) = \{q^6(q^2 - 1)^2, q^2 + q + 1, q^2 - q + 1\}$; if $3 \mid q - 1$, then $\text{OC}(G) = \{q^6(q^2 - 1)^2(q^2 + q + 1), q^2 - q + 1\}$; and if $3 \mid q + 1$, then $\text{OC}(G) = \{q^6(q^2 - 1)^2(q^2 - q + 1), q^2 + q + 1\}$. Thus if $3 \mid q$ and $M = G_2(q)$, then it has been proved that Thompson's conjecture holds in [4]. So it is necessary to discuss the case $3 \nmid q$.

LEMMA 10. *Let G be a finite group with $Z(G) = 1$ and $M = G_2(q)$, where $3 \mid q - 1$. If $N(G) = N(M)$, then $G \cong M$.*

Proof.

Step 1. Prove that G is neither a Frobenius nor a 2-Frobenius group.

(1) Prove that G is not a Frobenius group.

If G is a Frobenius group, then, by Lemma 3, $\text{OC}(G) = \{|H|, |K|\}$, where H and K are the Frobenius kernel and the Frobenius complement, respectively.

Suppose $2 \mid |H|$. Then $|K| = q^2 - q + 1$. Let p be a prime and $p \mid (q^2 + q + 1)/3$. Then $|S_p| < \frac{1}{3}(q^2 + q + 1) < q^2 - q + 1$. Since H is nilpotent, S_p is normal in G . Then $q^2 - q + 1 \mid |S_p|$ by Lemma 8, a contradiction.

Suppose $2 \mid |K|$. Then for any Sylow subgroup of H , we have $|S_p| < |K|$, contradicting $|K| \mid |S_p| - 1$ by Lemma 8.

(2) Prove that G is not a 2-Frobenius group.

If G is a 2-Frobenius group, then G has a normal series $H \trianglelefteq K \trianglelefteq G$ such that $|K/H| = q^2 - q + 1$ by Lemma 4. Since $q \equiv 1 \pmod{3}$, $q \geq 4$.

If q is odd, then $(q+1)/2$ or $(q-1)/2$ is an odd integer > 2 . When $(q-1)/2$ is odd, we have $((q-1)/2)^2(q^2 + q + 1)/3 \geq 3(q^2 + q + 1) > |K/H|$. By Lemma 4, $|G/K| < q^2 - q + 1$. Thus there exists a prime p

such that $p \mid ((q-1)/2)^2(q^2+q+1)/3$, $p \mid |H|$. We shall encounter a contradiction by similar reasoning as in (1). And we can deal with the case $(q+1)/2$ odd similarly.

If q is even, then $(q-1)^2(q^2+q+1)/3 > q^2 - q + 1$. Taking a prime p divides $(q-1)^2(q^2+q+1)/3$, similar to (1) we also encounter a contradiction. So (2) and then Step 1 follow.

By Step 1 and Lemma 7 we know that G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$, such that K/H is a simple group and $t(K/H) \geq 2$. Therefore we must check each possibility of K/H . By the classification theorem of finite simple groups and [6, 8], we may list all the possibilities of K/H as shown in Table I.

Now we must check all possibilities of K/H one by one. Since the procedure of checking is the same and trivial, we show several typical cases only.

Step 2. Prove that K/H is not an alternating group A_n .

Suppose $K/H = A_n$. Since $t(K/H) \geq 2$, we have that $n = p$, $p+1$, or $p+2$ by Table I, where p is an odd prime ≥ 5 .

$$\text{OC}(K/H) = \begin{cases} \{p\} & n, n-2 \text{ not all primes} \\ \{p, p-2\} & n, n-2 \text{ primes} \end{cases}$$

By Lemma 7 we have $q^2 - q + 1 = p$ or $p-2$. If $q^2 - q + 1 = p$, then $q^2 - q - 1 \mid |G|$ since $p-1 \mid |K/H|$. But it is easy to check that $(|G|, q^2 - q - 1) = 1$, a contradiction. If $q^2 - q + 1 = p-2$, then $p = q^2 - q + 3$, which implies $q^2 - q + 2 \mid (q^2 + q + 1)/3$ for $p \mid |G|$, a contradiction.

Step 3. Prove that K/H cannot be a simple group of type $A_n(q')$ or ${}^2A_n(q')$.

This step is the most difficult one; the method used here is suitable to other steps.

TABLE I

| | |
|---|---------------------------|
| $A_n(q)$, $(n, q) = (1, q)$ or $(p-1, q)$ or (p, q) and $q-1 \mid p-1$ | $G_2(q)$, $q > 2$ |
| ${}^2D_n(q)$, $(n, q) = (p+1, 2)$ or $(2^m, q)$ or $(p, 3)$, $p \geq 5$ | ${}^2G_2(q)$ |
| $E_7(2)$, $E_7(3)$ | ${}^2B_2(q)$ |
| A_n , $n = 5, 6$ or $n = p, p+1, p+2$ | ${}^2F_4(2)'$ |
| $B_n(q)$, $n = 2^m$, $m \geq 2$ | $B_p(3)$, $q = 2, 3$ |
| $C_n(q)$, $n = 2^m$, $m \geq 1$ | $C_p(q)$, $q = 2, 3$ |
| $D_p(q)$, $p \geq 5$, $q = 3, 5$ | $D_{p+1}(3)$, $p \geq 3$ |
| $E_6(q)$ | $F_4(q)$, $q > 2$ |
| ${}^2A_n(q)$, $(n, q) = (p-1, q)$ or (p, q) and $q+1 \mid p+1$ | $E_8(q)$ |
| ${}^3D_4(q)$ | ${}^2E_6(q)$ |
| A sporadic simple group | $G_2(2)'$ |

First we prove that K/H cannot be $A_n(q')$.

If $K/H = A_n(q')$, then $(n, q') = (1, q')$ or $(p' - 1, q')$ or (p', q') and $q' - 1 \mid p' - 1$ by Table I, where p is an odd prime.

(1) Let $n = 1$ and $q' \equiv 1 \pmod{4}$. Then by [2, Table 2] the odd order components of K/H are q' and $(q' + 1)/2$. Thus $q^2 - q + 1 = q'$ or $(q' + 1)/2$ by Lemma 7. If $q^2 - q + 1 = q'$, then $|A_1(q')| = \frac{1}{2}q(q-1)(q^2 - q + 1)(q^2 - q + 2)$. Since $|A_1(q')| \mid |G|$, $(q^2 - q + 2)/2$ divides $q^6(q-1)^2(q+1)^2(q^2 + q + 1)(q^2 - q + 1)$ and so $(q^2 - q + 2)/2$ divides $(q^2 + q + 1)$ because $(q^2 - q + 2)/2$ is coprime to $q(q^2 - 1)$. But $q^2 + q + 1 = (q^2 - q + 2) + 2q - 1$. Then $(q^2 - q + 2)/2$ divides $2q - 1$. Because $q \equiv 1 \pmod{3}$, $q \geq 4$. It is easy to check that $q = 4$; otherwise $(q^2 - q + 2)/2 > 2q - 1$ if $q > 4$. Thus $|G| = 2^{12} \cdot 3^3 \cdot 5^2 \cdot 7 \cdot 13$ and $A_1(13) = 2^2 \cdot 3 \cdot 7 \cdot 13$. By Lemma 7, $|G/K| \mid 2$ and $5^2 \mid |H|$. Since H is nilpotent, G has a normal subgroup of order 25, which leads to $13 \mid 24$ by Lemma 8, which is impossible.

(2) Let $n = 1$ and $q' \equiv -1 \pmod{4}$. Then the two odd order components of K/H are q' and $(q' - 1)/2$. By Lemma 7 we have $q^2 - q + 1 = q'$ or $(q' - 1)/2$. We can use the same method as in (1) to deal with it if $q^2 - q + 1 = q'$. If $q^2 - q + 1 = (q' - 1)/2$, then $q' \equiv 1 \pmod{D(M)}$, and so $(q, q') \neq 1$ or $q = 4$ and $q' = 3^3$ by Lemma 9. If $(q', q) \neq 1$, that is, $q = r^k$ and $q' = r^s$ for some prime r , it is easy to show this is impossible. If $q = 4$ and $q' = 27$, then $|A_1(q')| = 2^3 \cdot 3^3 \cdot 7 \cdot 13$. By similar reasoning as in (1), we have $|G/K| \mid 12$ and $5^2 \mid |H|$, so that G has a normal subgroup of order 25, and then $13 \mid 24$ by Lemma 8, a contradiction.

(3) Let $n = 1$ and $4 \mid q'$. Then the odd order components of K/H are $q' + 1$ and $q' - 1$, so that $q^2 - q + 1 = q' + 1$ or $q' - 1$. Since q' is a power of 2, $q^2 - q + 1 \neq q' + 1$ and $q^2 - q + 1 = q' - 1$. Moreover $q' \equiv 1 \pmod{q^2 - q + 1}$, which means that $2 \mid q$ or $(q, q') = (4, 27)$ by Lemma 9. Obviously, it is incorrect that $(q, q') = (4, 27)$ for $4 \mid q'$. If $2 \mid q$, then $q = q' = 4$ by $q^2 - q + 1 = q' - 1$, and $|G| = 2^5 \cdot 3^3 \cdot 7$ and $|A_1(4)| = 4 \cdot 3 \cdot 5$, which is impossible by similar reasoning.

(4) Let $n = 2$ and $q' = 4$. Then K/H has odd order components 5, 7, and 9. Thus $q^2 - q + 1 = 5, 7$, or 9 . It is easy to show that $q = 2$ for $q \equiv 1 \pmod{3}$, a contradiction.

(5) Let $n = p - 1$ and $(n, q') \neq (2, 4)$ where p is an odd prime. Then K/H has only one odd order component $(q'^p - 1)/[(q' - 1)(p, q' - 1)]$ and so $(q'^p - 1)/[(q' - 1)(p, q' - 1)] = q^2 - q + 1$. Hence $q'^p \equiv 1 \pmod{q^2 - q + 1}$ and again $(q, q') \neq 1$ or $(q, q'^p) = (4, 27)$ by Lemma 7.

If $(q, q') \neq 1$ and $(p, q' - 1) = 1$, then $q'^{p-1} + \cdots + q' + 1 = q^2 - q + 1$ and

$$q'(q'^{p-2} + \cdots + q' + 1) = q(q - 1),$$

which is impossible.

If $(q, q') \neq 1$ and $(p, q' - 1) = p$, then $(q'^{p-1} + \cdots + q' + 1)/p = (q'^p - 1)/(q' - 1)(p, q' - 1) = q^2 - q + 1$. Therefore $p - 1 \equiv 0 \pmod{q}$ or $p - 1 \equiv 0 \pmod{q'}$. Obviously, the latter one is impossible for $p \mid q' - 1$. If the first one holds and $p \geq 5$, then $q^3 > (q'^p - 1)/(q' - 1)p > q'^3$. This means $q > q'$, contradicting $q \mid p - 1$ and $p \mid q' - 1$. If the first one holds and $p = 3$, then $q \mid 2$. Thus $q = 2$, a contradiction.

If $q = 4$ and $q'^p = 27$, then $q' = 3$, $p = 3$, $q^2 - q + 1 = 13$, $|G| = 2^{12} \cdot 3^3 \cdot 5^2 \cdot 7 \cdot 13$ and $|K/H| = 2^4 \cdot 3^3 \cdot 13$. By the same reasoning as in (2), we have $7 \parallel |H|$, a contradiction.

(6) Let $(n, q') = (p, q')$ and $q' - 1 \mid p + 1$. Then K/H has an odd order component $(q'^p - 1)/(q' - 1)$. Similar to (5), it can be shown that this case is also impossible.

Until now we have proved that K/H cannot be $A_n(q')$. By the same method we can prove that K/H cannot be a simple group of type ${}^2A_n(q')$ because the odd order components of ${}^2A_n(q')$ are of the same form of those of $A_n(q')$.

Step 4. Prove that K/H cannot be simple groups of type $B_n(q')$ and $C_n(q')$.

Since $B_n(q')$ and $C_n(q')$ have the same order and the same odd order components, it is sufficient to discuss one of them, for example, $B_n(q')$.

If $K/H = B_n(q')$, then $(n, q') = (2^m, q')$ for $m > 1$ or (p, k) with p a prime and $k = 2$ or 3 by Table I.

(1) If $(n, q') = (2^m, q')$ with $m > 1$, then K/H has an odd order component $(q'^n + 1)/(2, q' - 1)$. Therefore $(q'^n + 1)/(2, q' - 1) = q^2 - q + 1$, so that $q'^{n^2} > (q(q - 1))^n$. If $n \geq 8$, then $q'^{n^2} > q^6$, which is impossible by Lemma 9. Hence $n < 8$.

If $n = 4$, then $q'^{n^2} > q^4(q - 1)^4$, while $q > 2$, $q'^{n^2} > q^6$, a contradiction by Lemma 9.

If $n = 2$, then $q^2 - q + 1 = (q'^2 + 1)/(2, q' - 1)$. It is easy to show that $2 \nmid q'$ and $q'^4 > q^2(q - 1)^2 > q^3$. Then we have $(q, q') \neq 1$ by Lemma 9. It is easy to check that this is impossible.

(2) If $(n, q') = (p, k)$ for p a prime and $k = 2$ or 3 , then K/H has an odd order component $(k^p - 1)/(2, k - 1)$, which is equal to $q^2 - q + 1$. By the same reasoning as in (1), we can show this is also impossible. Step 4 follows.

Step 5. Prove that K/H cannot be a simple group of type $D_n(q')$.

If $K/H = D_n(q')$, then $(n, q') = (p, k)$ or $(p' + 1, 3)$, where p is a prime ≥ 5 , p' is a prime ≥ 3 , and $k = 2$ or 3 or 5 by Table I.

If $(n, q') = (p, k)$, p and k as above, then K/H has an order component $(k^p - 1)/(4, k - 1)$. Thus $(k^p - 1)/(4, k - 1) = q^2 - q + 1$ and so $k^p \equiv 1 \pmod{q^2 - q + 1}$ and $k^{p(p-1)} > q^4(q-1)^4$. Then, by Lemma 7, since $q \geq 4$, $k^{p(p-1)} > q^6$ for $q > 2$, a contradiction.

If $(n, q') = (p' + 1, 3)$, then $D_n(q')$ has an odd order component $(3^{p'} - 1)/2$. For $p' \geq 5$, by the same process as in (1), it can be proved that K/H cannot be $D_n(q')$. If $p' = 3$, then $q^2 - q + 1 = 13$ and so $q = 3$ or 4 . But by $q \equiv 1 \pmod{3}$ we have $q = 4$, and so that $3^{12} \mid |K/H|$, contradicting $3^{12} \nmid |G|$. Step 5 follows.

Step 6. Prove that K/H cannot be a simple group of types: ${}^2B_2(q')$, ${}^2D_n(q')$, $E_6(q')$, ${}^2E_6(q')$, $F_4(q')$, ${}^2F_4(q')$, ${}^2G_2(q')$, and ${}^3D_4(q')$.

Since the processes to prove that K/H cannot be a simple group of the types listed above are the same as in those in Steps 3, 4, and 5, we omitted the proofs here.

Step 7. Prove that K/H cannot be $E_8(q')$.

By [4, Table 1], the odd order components of $E_8(q')$ are all $\leq q'^9$. If $K/H = E_8(q')$, then $q^2 - q + 1$ equals one of the odd order components of $E_8(q')$. Therefore $q'^9 > q$ and $q'^{120} > q^{10}$, which is impossible by $q'^{120} \mid |E_8(q')|$ and Lemma 7. Step 7 follows.

Step 8. Prove that K/H cannot be any one of sporadic simple groups, $E_7(2)$, $E_7(3)$, and ${}^2F_4(2)$.

By Tables 2–4 in [2] any odd order component of any simple group listed above is one of the following primes: 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, 59, 67, 71, 73, 127, 757, 1093. Hence $q^2 - q + 1$ equals one of these primes. Then $q = 4$ by solving these equations, so that $|G| = |G_2(4)| = 2^{12} \cdot 3^3 \cdot 5^2 \cdot 7 \cdot 13$, which leads to $\text{Max } \pi(K/H) = \text{Max } \pi(G) = 13$. Then $K/H = Sz$ by [8]. But $11 \mid |Sz|$ and $11 \nmid |G|$, a contradiction.

Step 9. Prove that $G \cong M$.

From Steps 1–9, we know that $K/H = G_2(q')$. If $q' \equiv -1 \pmod{3}$, then K/H has odd order component $q'^2 + q' + 1$, so $q'^2 + q' + 1 = q^2 - q + 1$ and $q'(q' + 1) = q(q - 1)$. Hence $q' = q - 1$ and $q' \equiv 0 \pmod{3}$ for $q \equiv 1 \pmod{3}$, a contradiction to $q' \equiv -1 \pmod{3}$.

If $q' \equiv 0 \pmod{3}$, then K/H has two odd order components $q'^2 + q' + 1$ and $q'^2 - q' + 1$. Thus $q^2 - q + 1 = q'^2 + q' + 1$ or $q'^2 - q' + 1$, which implies $q = q' + 1$ or $q = q'$. Since $q' \equiv 0 \pmod{3}$ and $q \equiv 1 \pmod{3}$, $q = q' + 1$ and $q'^6 = (q - 1)^6$. Therefore $q'^6 > q^3$ for $q \geq 4$. Then $(q', q) \neq 1$ by Lemma 7, a contradiction.

If $q' \equiv 1 \pmod{3}$, then $q'^2 - q' + 1 = q^2 - q + 1$, so that $q' = q$ and $G = K/H \cong M$.

This is the end of the proof of Lemma 9.

The proof of the following lemma is similar to that of Lemma 10.

LEMMA 11. *Let G be a finite group with $Z(G) = 1$ and $M = G_2(q)$, where $3 \mid q + 1$. If $N(G) = N(M)$, then $G \cong M$.*

LEMMA 12. *Let G be a finite group with $Z(G) = 1$ and $M = G_2(2)$ or $G_2(2)'$. If $N(G) = N(M)$, then $G \cong M$.*

Proof. First, since $G_2(2)' = {}^2A_2(3)$, it has been discussed in Lemma 10.

Second, by [5, Page 14], we know that the prime graph of $G_2(2)$ has two components: $2^6 \cdot 3^3$ and 7. Then if $M = G_2(2)$, we have to repeat the processes used in the proof of Lemma 10 by Lemmas 1, 2, and 5. Similar to Step 1 in the proof of Lemma 10, we can prove that G cannot be a Frobenius group or a 2-Frobenius group. Therefore G has a normal series $H \trianglelefteq K \trianglelefteq G$, such that K/H is a simple group with $t(K/H) \geq 2$. Since $|K/H| \mid 2^6 \cdot 3^3 \cdot 7$, K/H may be one of following simple groups by [5]: $L_2(3)$ ($2^3 \cdot 3 \cdot 7$), $L_2(8)$ ($2^3 \cdot 3^2 \cdot 7$), and $G_2(2)'$.

If $K/H = L_2(7)$, then G/K has an order of a factor of $|\text{Out}(L_2(7))|$, that is, $|G/K| \mid 2$. Hence H has a 3-Sylow subgroup of order 9 that is normal in G , a contradiction by Lemma 8. So $K/H \neq L_2(7)$.

If $K/H = L_2(8)$, then $|G/K| \mid 3$ by Lemma 5. Therefore the 2-Sylow subgroup N of H is of order 8. by Lemma 8, N has exactly two conjugacy classes, one containing seven elements and one element. That is to say, for any $x \in N$ and $x \neq 1$, we have $|\text{Cl}_G(x)| = 7$, contradicting $7 \notin N(G_2(2))$ by [5, p. 13]. Hence $K/G \neq L_2(8)$.

If $K/H = G_2(2)'$, then $|H| = 1$ or 2. By Lemma 8, $|H| = 1$ and $|G/K| = 2$, which leads to $G \cong M$.

Proof of the Main Theorem. The theorem follows from Lemmas 10–12.

REFERENCES

1. G. Y. Chen, On Thompson's conjecture—for sporadic groups (Chinese), in "Proceedings of the China Association of Science and Technology, First Academic Annual Meeting of Youths," pp. 1–6, Chinese Science and Technology Press, Beijing, 1992.
2. G. Y. Chen, On the structure of Frobenius group and 2-Frobenius group, *J. Southwest China Normal Univ.* **20**, No. 5 (1995), 485–487.
3. G. Y. Chen, A new characterization of sporadic simple groups, *Algebra Colloq.* **3**, No. 1 (1996), 49–58.

4. G. Y. Chen, On Thompson's conjecture, *J. Algebra* **185** (1996), 184–193.
5. J. H. Conway, R. T. Curtis, *et al.*, “Atlas of Finite Groups,” Clarendon, Oxford, 1985.
6. A. S. Kondratev, Prime graph components of finite simple groups, *Mat. USSR Sb.* **67**, No. 1 (1990), 235–247.
7. W. J. Shi and J. X. Bi, A characteristic property of each finite projective special group, in “Lecture Notes in Mathematics,” Vol. 1456, pp. 171–180, Springer-Verlag, Berlin/New York, 1990.
8. J. S. Williams, Prime graph components of finite groups, *J. Algebra* **69** (1981), 487–513.